

#11502

AKT1 (phospho-Thr450) Antibody

Catalog: #11502-1 50µl **Orders:** order@signalwayantibody.com
 #11502-2 100µl **Support:** tech@signalwayantibody.com
Storage: Store at -20°C/1 year **Web:** www.signalwayantibody.com



Application	Species Reactivity	Source	Molecular Wt.
WB IHC IF	Human Mouse Rat	Rabbit Polyclonal Ab	60KD

Description: Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.

Specificity: The antibody detects endogenous level of AKT1 only when phosphorylated at threonine 450.

Immunogen: Peptide sequence around phosphorylation site of threonine 450 (T-I-T(p)-P-P) derived from Human AKT1.

Formulation: Supplied at 1.0mg/mL in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.


Synonyms: RAC-PK-alpha
Protein kinase B

Accession No.: Swiss-Prot#: P31749 NCBI Gene#: 207
NCBI Protein#: NP_001014431.1

Background: General protein kinase capable of phosphorylating several known proteins. Phosphorylates TBC1D4. Signals downstream of phosphatidylinositol 3-kinase (PI3K) to mediate the effects of various growth factors such as platelet-derived growth factor (PDGF), epidermal growth factor (EGF), insulin and insulin-like growth factor I (IGF-I). Plays a role in glucose transport by mediating insulin-induced translocation of the GLUT4 glucose transporter to the cell surface. Mediates the antiapoptotic effects of IGF-I. Mediates insulin-stimulated protein synthesis by phosphorylating TSC2 at 'Ser-939' and 'Thr-1462', thereby activating mTORC1 signaling and leading to both phosphorylation of 4E-BP1 and in activation of RPS6KB1. Promotes glycogen synthesis by mediating the insulin-induced activation of glycogen synthase.

References:

Xing J, et al. (1998) Mol Cell Biol 18(4): 1946-55.
 Tan Y, et al. (1996) EMBO J; 15(17): 4629-42.
 Hao, M. et al. (1996) J. Biol. Chem. 271, 29380-29385.
 Mayo LD, et al. (2001) Biol Chem; 276(27): 25184-9.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using AKT1 (Phospho-Thr450) Antibody #11502 (left) or the same antibody preincubated with blocking peptide (right).


Citation: If you publish research using #11502 please [let us know](#).

Related Pathway: Akt, Insulin/Glucose, Jak/Stat, Cytoskeletal/Adhesion, Wnt/beta-catenin


Note: For western blotting, incubate membrane with diluted antibody in 5% nonfat milk, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Recommended Dilutions:

Western blotting 1:500~1:1000
 Immunohistochemistry 1:50~1:100
 Immunofluorescence 1:100~1:200



Immunofluorescence staining of methanol-fixed HeLa cells using AKT1 (phospho-Thr450) Antibody #11502.



Western blot analysis of extracts from 293 cells untreated or treated with EGF using AKT1 (phospho-Thr450) Antibody #11502.